

Quantitation of Changes in Protein Expression between Chicken Dark and White Meat by PMF and MSMS.

Kenneth C. Parker, Stephen J. Hattan, Marvin Vestal. *SimulTof Corporation, Sudbury, MA*

WP20 #411

Introduction

Many multicellular organisms contain several different kinds of muscle fibers, which result from the expression of distinct combinations of major muscle protein genes. Muscles are classifiable as fast or slow, based predominantly on myosin heavy chain genes. Many other major muscle proteins derive from a family of homologous genes. In humans, myosin heavy chain usage changes with disease, with muscle type and according to exercise patterns. Moreover, muscle preparations tend to contain a mixed population of slow twitch and fast twitch muscle. Chicken was chosen as a model organism because meat is readily available enriched in slow fibers (dark meat) or fast fibers (white meat). We have determined how major muscle proteins differ across meat types.

Methods

Meat samples were homogenized, reduced, alkylated, and digested with trypsin. Digests were subjected directly to PMF, resulting in the identification of tens of proteins. In addition, the same digest preparations were subjected to HPLC-MALDI analysis, and parent MS spectra and MS-MS spectra were acquired. Each separation resulted in a peak list containing ~ 10000 peptide masses, measured with < 5 ppm accuracy with the use of two internal standards. PMF software was used that takes into account preferential arginine detection, and novel MS-MS software was developed to interpret high energy MS-MS spectra. The intensities from the same peak lists allowed quantitation of changes between muscle types. In the last few weeks, MS-MS spectra were gathered directly using SimulTof 300 Tandem.



14 samples:
6 white meat
8 dark meat


100 Most intense masses from each

Merged mass list with 226 masses and intensities

Workflows:

1. Perform PMF on tryptic digest of whole tissue.
2. Perform PCA on mass lists
3. Acquire MS-MS spectra using LC-MALDI
4. Collect MS-MS spectra directly from unseparated digests

Fig. 1. PCA analysis




3 distinct categories

Fig1B,C. Mass-based PCA

MYH: masses mapped to any myosin heavy chain isoform. In mass-based PCA, the strongest drivers correspond to MYH, though some MYH masses (in black in Fig.1B) are constant. Most glycolytic enzyme masses are strongest in white meat, while creatine kinase (CK) is enriched in both dark meat categories. Heat shock protein HSPB2 is enriched in the dark meat category 'd1'.

Fig 2. Distribution of selected masses over the 14 samples



Amino acids that are variable between myosin isoforms are color-coded as on Fig. 1. Some peptides have only 2 forms. Many other MYH peptides are shared between these isoforms. Peptides aligned to one MYH isoform, with 1st aa listed.

Fig. 3. Same region following LC separation. High energy msms spectra.




Fig. 5. MS-MS spectrum of SYELPDGQVITIGNER from actin (strongest peak) from whole digest using SimulTof 300 tandem. Mascot Score 104.





Fig. 4. Comparison of MS and MS-MS spectra obtained via LC-MALDI vs spectra obtained directly from separated digest focusing on mass 1869. 8.



Conclusions:

- Using MS, can see differences between meat samples.
- Using PCA, one can separate samples into 3 categories.
 - White meat, two separate dark meat categories.
- Can identify 5-10 proteins from chicken muscle by PMF.
- Confirmation of PMF by MSMS can be performed directly on unseparated digests, resulting in ~15 protein IDs from whole chicken muscle so far.
- More confirmation obtained by MSMS following LC-MALDI.
- Differences correspond to MYH isoforms, glycolytic enzymes, etc.

References:

- 1.) Parker KC. Scoring Methods in MALDI Peptide Mass Fingerprinting: ChemScore and the ChemApplix Program. JASMS 2002;13:22-39.

Funding: SBIR grant 1R44GM090389-01A1.

